ELSEVIER

Biomembranes

Vol. 1415, 1999

BIOCHIMICA ET BIOPHYSICA ACTA

BBA

Author Index

Altendorf, K., see Heermann, R. (1415) 114
Altendorf, K., see Gaßel, M. (1415) 77
Auger, M., see Bouchard, M. (1415) 181

Benning, C., see Härtel, H. (1415) 205
Bouchard, M., Le Guernevé, C. and Auger, M.
Comparison between the dynamics of lipid/gramicidin A systems in the lamellar and hexagonal phases: a solid-state ^{13}C NMR study (1415) 181
Burson, K.K., see Hall, M.P. (1415) 101

Cabral-Lilly, D., see Li, X. (1415) 23
Cailler, F., Howell, S. and Crine, P.
Role of the glycosyl-phosphatidylinositol anchor in the intracellular transport of a transmembrane protein in Madin-Darby canine kidney cells (1415) 1
Cajal, Y., see Oh, J.-T. (1415) 235
Chan, H.C., see So, S.C. (1415) 261
Cheng, C.Y., see So, S.C. (1415) 261
Cheng, Y., see Hiroaki, Y. (1415) 361
Chung, Y.W., see So, S.C. (1415) 261
Crine, P., see Cailler, F. (1415) 1
Cruz, A., Marsh, D. and Pérez-Gil, J.
Rotational dynamics of spin-labelled surfactant-associated proteins SP-B and SP-C in dipalmitoylphosphatidylcholine and dipalmitoylphosphatidylglycerol bilayers (1415) 125

de Kruijff, B., see Harris, F. (1415) 10
Demel, R., see Harris, F. (1415) 10
Deslauriers, R., see Gruwel, M.L.H. (1415) 41
Dhurjati, P.S., see Oh, J.-T. (1415) 235
Duclohier, H., Snook, C.F. and Wallace, B.A.
Antiamoebin can function as a carrier or as a pore-forming peptaibol (1415) 255

Eckstein-Ludwig, U., see Liu, Y. (1415) 246
Epstein, W., see Gaßel, M. (1415) 77
Essigmann, B., see Härtel, H. (1415) 205

Fei, J., see Liu, Y. (1415) 246
Franzin, C.M., Macdonald, P.M., Polozova, A. and Winnik, F.M.
Destabilization of cationic lipid vesicles by an anionic hydrophobically modified poly(*N*-isopropylacrylamide) copolymer: a solid-state ^{31}P NMR and ^{2}H NMR study (1415) 219
Fromm, M.F., Leake, B., Roden, D.M., Wilkinson, G.R. and Kim, R.B.
Human MRP3 transporter: identification of the 5′-flanking region, genomic organization and alternative splice variants (1415) 369
Fujiyoshi, Y., see Hiroaki, Y. (1415) 361

Gaßel, M., Siebers, A., Epstein, W. and Altendorf, K.
Assembly of the Kdp complex, the multi-subunit K^+-transport ATPase of *Escherichia coli* (1415) 77
Gioioso, S., see Moore, D.J. (1415) 342
Grima, J., see So, S.C. (1415) 261
Gruner, S.M., see Li, X. (1415) 23
Gruwel, M.L.H., Kuzio, B., Xiang, B., Deslauriers, R. and Kupriyanov, V.V.
Temperature dependence of monovalent cation fluxes in isolated rat hearts: a magnetic resonance study (1415) 41

Hall, M.P., Burson, K.K. and Huestis, W.H.
Interactions of a vesicular stomatitis virus G protein fragment with phosphatidylserine: NMR and fluorescence studies (1415) 101
Harauz, G., see Palaniyar, N. (1415) 85
Harris, F., Demel, R., de Kruijff, B. and Phoenix, D.A.
An investigation into the lipid interactions of peptides corresponding to the C-terminal anchoring domains of *Escherichia coli* penicillin-binding proteins 4, 5 and 6 (1415) 10
Härtel, H., Essigmann, B., Lokstein, H., Hoffmann-Benning, S., Peters-Kottig, M. and Benning, C.
The phospholipid-deficient *pho1* mutant of *Arabidopsis thaliana* is affected in the organization, but not in the light acclimation, of the thylakoid membrane (1415) 205
Heermann, R., Altendorf, K. and Jung, K.
The turgor sensor KdpD of *Escherichia coli* is a homodimer (1415) 114
Heimburg, T.
Mechanical aspects of membrane thermodynamics. Estimation of the mechanical properties of lipid membranes close to the chain melting transition from calorimetry (1415) 147
Hernández-Hernández, A., see Rocio Marcos, M. (1415) 163
Hiroaki, Y., Mitsuoka, K., Cheng, Y., Hiroaki, H. and Fujiyoshi, Y.
Influence of various nucleotides on the in situ crystallization of Ca^{2+}-ATPase (1415) 361
Hiroaki, H., see Hiroaki, Y. (1415) 361
Hirsh, D.J., see Li, X. (1415) 23
Hoffmann-Benning, S., see Härtel, H. (1415) 205
Howell, S., see Cailler, F. (1415) 1
Huang, S.-G., see Klingenberg, M. (1415) 271
Huestis, W.H., see Hall, M.P. (1415) 101
Hui, S.W., see Langner, M. (1415) 323
Hunter, G.W. and Squier, T.C.
Phospholipid acyl chain rotational dynamics are independent of headgroup structure in unilamellar vesicles containing binary mixtures of dioleoyl-phosphatidylcholine and dioleoyl-phosphatidylethanolamine (1415) 63

Ikigai, H., Ono, T., Nakae, T., Otsuru, H. and Shimamura, T.
Two forms of *Vibrio cholerae* O1 El Tor hemolysin derived from identical precursor protein (1415) 297
Ivanov, I.T.
Low pH-induced hemolysis of erythrocytes is related to the entry of the acid into cytosole and oxidative stress on cellular membranes (1415) 349

Jack, D.L., see Young, G.B. (1415) 306
Jain, M.K., see Oh, J.-T. (1415) 235
Janoff, A.S., see Li, X. (1415) 23
Jo, O.D., see Yanagawa, N. (1415) 56
Jung, K., see Heermann, R. (1415) 114

Kim, R.B., see Fromm, M.F. (1415) 369
Klingenberg, M. and Huang, S.-G.
Structure and function of the uncoupling protein from brown adipose tissue (1415) 271
Kupriyanov, V.V., see Gruwel, M.L.H. (1415) 41
Kuzio, B., see Gruwel, M.L.H. (1415) 41

Langner, M. and Hui, S.W.
Merocyanine 540 as a fluorescence indicator for molecular packing stress at the onset of lamellar-hexagonal transition of phosphatidylethanolamine bilayers (1415) 323
Le Guernevé, C., see Bouchard, M. (1415) 181
Leake, B., see Fromm, M.F. (1415) 369
Leung, P.S., see So, S.C. (1415) 261
Li, X., Hirsh, D.J., Cabral-Lilly, D., Zirkel, A., Gruner, S.M., Janoff, A.S. and Perkins, W.R.
Doxorubicin physical state in solution and inside liposomes loaded via a pH gradient (1415) 23
Liu, Y., Eckstein-Ludwig, U., Fei, J. and Schwarz, W.
Effect of mutation of glycosylation sites on the Na^+ dependence of steady-state and transient currents generated by the neuronal GABA transporter (1415) 246
Llanillo, M., see Rocío Marcos, M. (1415) 163
Lokstein, H., see Härtel, H. (1415) 205

Macdonald, P.M., see Franzin, C.M. (1415) 219
Marsh, D., see Cruz, A. (1415) 125
Massey, J.B.
Effect of cholesteryl hemisuccinate on the interfacial properties of phosphatidylcholine bilayers (1415) 193
Mendelsohn, R., see Moore, D.J. (1415) 342
Mitsuoka, K., see Hiroaki, Y. (1415) 361
Moore, D.J., Gioioso, S., Sills, R.H. and Mendelsohn, R.
Some relationships between membrane phospholipid domains, conformational order, and cell shape in intact human erythrocytes (1415) 342
Moscarello, M.A., see Palaniyar, N. (1415) 85

Nakae, T., see Ikigai, H. (1415) 297

Oh, J.-T., Cajal, Y., Dhurjati, P.S., Van Dyk, T.K. and Jain, M.K.
Cecropins induce the hyperosmotic stress response in *Escherichia coli* (1415) 235
Ono, T., see Ikigai, H. (1415) 297
Otsuru, H., see Ikigai, H. (1415) 297

Pajor, A.M.
Sequence of a pyrimidine-selective Na^+/nucleoside cotransporter from pig kidney, pkCNT1 (1415) 266
Palaniyar, N., Semotok, J.L., Wood, D.D., Moscarello, M.A. and Harauz, G.
Human proteolipid protein (PLP) mediates winding and adhesion of phospholipid membranes but prevents their fusion (1415) 85
Pérez-Gil, J., see Cruz, A. (1415) 125
Perkins, W.R., see Li, X. (1415) 23
Peters-Kottig, M., see Härtel, H. (1415) 205
Pflugmacher, D. and Sandermann Jr., H.
The lipid/protein interface as a target site for general anesthetics: a multiple-site kinetic analysis of synaptosomal Ca^{2+}-ATPase (1415) 174
Phoenix, D.A., see Harris, F. (1415) 10
Polozova, A., see Franzin, C.M. (1415) 219

Rocío Marcos, M., Sánchez-Yagüe, J., Hernández-Hernández, A. and Llanillo, M.
Amphiphilic and hydrophilic forms of acetylcholinesterase from sheep platelets (1415) 163
Roden, D.M., see Fromm, M.F. (1415) 369
Ruponen, M., Ylä-Herttuala, S. and Urtti, A.
Interactions of polymeric and liposomal gene delivery systems with extracellular glycosaminoglycans: physicochemical and transfection studies (1415) 331

Said, H.M., see Yanagawa, N. (1415) 56
Saier Jr., M.H., see Young, G.B. (1415) 306
Sánchez-Yagüe, J., see Rocío Marcos, M. (1415) 163
Sandermann Jr., H., see Pflugmacher, D. (1415) 174
Schote, U. and Seelig, J.
Interaction of the neuronal marker dye FM1-43 with lipid membranes (1415) 135
Schwarz, W., see Liu, Y. (1415) 246
Seelig, J., see Schote, U. (1415) 135
Semotok, J.L., see Palaniyar, N. (1415) 85
Shimamura, T., see Ikigai, H. (1415) 297
Siebers, A., see Gaßel, M. (1415) 77
Sills, R.H., see Moore, D.J. (1415) 342
Smith, D.W., see Young, G.B. (1415) 306
Snook, C.F., see Duclohier, H. (1415) 255
So, S.C., Wu, W.L., Grima, J., Leung, P.S., Chung, Y.W., Cheng, C.Y., Wong, P.Y.D., Yan, Y.C. and Chan, H.C.
Functional expression of sperm angiotensin II type I receptor in *Xenopus* oocyte: modulation of a sperm Ca^{2+}-activated K^+ channel (1415) 261
Squier, T.C., see Hunter, G.W. (1415) 63

Urtti, A., see Ruponen, M. (1415) 331

Van Dyk, T.K., see Oh, J.-T. (1415) 235

Wallace, B.A., see Duclohier, H. (1415) 255
Wilkinson, G.R., see Fromm, M.F. (1415) 369
Winnik, F.M., see Franzin, C.M. (1415) 219
Wong, P.Y.D., see So, S.C. (1415) 261
Wood, D.D., see Palaniyar, N. (1415) 85
Wu, W.L., see So, S.C. (1415) 261

Xiang, B., see Gruwel, M.L.H. (1415) 41

Yan, Y.C., see So, S.C. (1415) 261
Yanagawa, N., Jo, O.D. and Said, H.M.
Riboflavin transport by rabbit renal basolateral membrane vesicles (1415) 56
Ylä-Herttuala, S., see Ruponen, M. (1415) 331
Young, G.B., Jack, D.L., Smith, D.W. and Saier Jr., M.H.
The amino acid/auxin:proton symport permease family (1415) 306

Zirkel, A., see Li, X. (1415) 23